

Mg²⁺ ions mediate the interaction of intrinsically disordered nascent chains with the ribosome: implications for protein folding and aggregation in the early stages of protein life

Valeria Guzman-Luna, Silvia Cavagnero*

Department of Chemistry, University
of Wisconsin-Madison, 1101 University
Ave., Madison, Wisconsin, 53706, USA

*Author for correspondence:
Email: cavagnero@chem.wisc.edu

Received date: January 13, 2021
Accepted date: March 28, 2021

Copyright: © 2022 Guzman-Luna V,
et al. This is an open-access article
distributed under the terms of the
Creative Commons Attribution License,
which permits unrestricted use,
distribution, and reproduction in any
medium, provided the original author
and source are credited.

Citation: Guzman-Luna V, Cavagnero
S. Mg²⁺ ions mediate the interaction of
intrinsically disordered nascent chains
with the ribosome: implications for
protein folding and aggregation in the
early stages of protein life. Neurosci
Chron. 2022;3(1):3-5.

During the last few decades, the ribosome has been regarded primarily as a major cell player devoted to the catalysis of protein biosynthesis during translation [1-5]. It is therefore not surprising that several processes related to translation exploit the ribosome as a central hub. For instance, it is well-known that many events related to translational regulation are mediated by interactions between the ribosome and initiation, elongation or termination factors [6-9]. In addition, the ribosome is involved in mRNA-code recognition and proofreading [10-12] as well as in the control of translation rates via interactions with mRNA codons bearing high- and low-frequency [13-15] and associated with variable tRNA abundance within the translation machinery [16-18]. Interestingly, the ribosome also assists *de novo* protein structure formation by minimizing cotranslational aggregation, thus increasing the yield of native-protein production [19,20]. The latter event, however, has not been shown to require -- or even involve -- direct interactions between the ribosome and the nascent protein chain. A notable exception is that of nascent chains bearing either N-terminal signal sequences or translational-arrest tags. These proteins are known to establish short- or long-term contacts with various regions of the ribosome during translation [21-25]. In summary, until recently very little knowledge has been available about direct contacts between the ribosome and nascent polypeptides and proteins that do not carry signal or arrest sequences. Studies based on fluorescence depolarization in the frequency domain [26] and NMR spectroscopy [27-30] provided interesting data that are consistent with, but do not unequivocally establish, the presence of these interactions.

A recent investigation by Guzman-Luna *et al.* [31] added new knowledge to the field by showing that two bacterial ribosomal proteins, L23 and L29, interact with ribosome-bound nascent protein chains (RNCs) encoding an intrinsically disordered protein (IDP) and several of its mutants bearing variable net charge. This study is based on site-specific fluorescence labeling of RNCs at their N terminus, in combination with chemical crosslinking and Western blotting. Intriguingly, short RNCs were found to interact only with the L23 ribosomal protein while longer RNCs also weakly interact with an additional ribosomal protein, L29. The interacting proteins are located within a specific region of the large ribosomal subunit, which comprises the vestibule of the exit tunnel as well as the immediately adjacent outer region of the ribosome.

Importantly, a large fraction of the detected interactions is mediated by Mg²⁺ ions, and the extent of the contacts can be tuned up and down depending on the Mg²⁺ concentration in the medium [31]. Interestingly, Mg²⁺ ions have been long known to be ribosome-associated [32,33] and to play a key role in preserving the structural integrity of the ribosome [32,33]. However, no Mg²⁺-mediated processes involving nascent proteins were ever detected before the work by Guzman-Luna *et al.* In summary, the study by Guzman-Luna *et al.* [31] highlights a new role for the ribosome, which is found to interact with intrinsically disordered nascent chains in a Mg²⁺-mediated fashion.

On the other hand, some (ca 50%) of the interactions of nascent proteins with the ribosome were found to be Mg²⁺-independent. Based on a quantitative electrostatic mapping of the ribosomal surface, which contains a large nonpolar patch on the solvent-exposed region of L23 within the exit region of the ribosomal tunnel [31], it is possible that the Mg²⁺-independent interactions are dominated by the hydrophobic effect [34]. Some classical electrostatic interactions not involving Mg²⁺ may also be present. However, these types of interactions are surprisingly not dominant. Ongoing and future studies will shed additional light on the remaining unanswered questions regarding the detailed nature of these IDP-ribosomal-protein interactions.

The study by Guzman-Luna and coworkers [31] nicely complements the current knowledge on the effect of Mg²⁺ ions on non-ribosome-related functions. For instance, Mg²⁺ is known to play a seminal role in bacterial thermodynamic balance [35-37] and in proteostasis, including modulating the activity of the Hsc62 [38] and Hsp70 [39] molecular chaperones and altering the specificity of the Lon and ClpAP proteases [40].

A few major lingering questions are (a) whether the above interactions are also present in the case of proteins with a foldable sequence, as opposed to IDPs, (b) what the role of the Mg²⁺-dependent and Mg²⁺-independent ribosome-RNC interactions is, and (c) whether similar interactions are also present across eukaryotic organisms. Ongoing research in the Cavagnero group (Guzman-Luna *et al.*, manuscript in preparation) suggests that extensive ribosome-RNC interactions are also present in the case of foldable protein sequences. However, their ion dependence is different and much more complex.

On a different note, while analogous studies on eukaryotic organisms are not available yet, it is notable that intracellular Mg²⁺ concentration is known to be important for human brain cytopathies [41], and that the concentration of Mg²⁺ ions has an effect on the extent of post-translational protein aggregation, as recently found in the case of TDP-43 [42]. Further, several known proteinopathies are related to the misfolding and aggregation of intrinsically disordered polypeptides and proteins [43,44]. It is also worth noting that the prokaryotic and eukaryotic ribosomes share a similar structure and overall highly negative electrostatic surface potential [45]. Incidentally, the ribosome (or fragments thereof) has been previously proposed to serve as a chaperone [46,47]. Therefore, despite the undeniable differences between ribosomes in all classes of organisms [6,48], it is tempting to hypothesize that Mg²⁺-mediated interactions between intrinsically disordered nascent proteins and the ribosome may play a role in regulating protein quality in bacteria and perhaps even in higher organisms. The known Mg²⁺ dependence of translation efficiency is consistent with this hypothesis [37].

In conclusion, Mg²⁺ mediates the interaction of intrinsically disordered nascent proteins and the bacterial ribosome [31]. Future studies will reveal whether Mg²⁺-mediated RNC-ribosome contacts influence the quality of *de-novo* synthesized proteins in bacteria. In the case of higher organisms, if confirmed to be present, ribosome-RNC interactions may influence the course of diseases that critically depend on the *de novo* generation of proteins devoid of aggregation. While more studies clearly need to be carried out to unequivocally identify additional trends, the present findings promise exciting opportunities for future discovery.

Acknowledgments

Funding from the National Science Foundation (MCB-2124672 and CBET-1912259) is gratefully acknowledged.

References

1. Davidovich C, Belousoff M, Bashan A, Yonath A. The evolving ribosome: from non-coded peptide bond formation to sophisticated translation machinery. *Research in Microbiology.* 2009;160:487-492.
2. Erlacher MD, Polacek N. Ribosomal catalysis - The evolution of mechanistic concepts for peptide bond formation and peptidyl-tRNA hydrolysis. *RNA Biology.* 2008; 5:5-12.
3. Rodnina MV, Beringer M, Wintermeyer W. How ribosomes make peptide bonds. *Trends in Biochemical Sciences.* 2007;32:20-26.
4. Aitken CE, Petrov A, Puglisi JD. Single Ribosome Dynamics and the Mechanism of Translation. *Annual Review of Biophysics.* 2010;39:491-513.
5. Nissen P, Hansen J, Ban N, Moore PB, Steitz TA. The structural basis of ribosome activity in peptide bond synthesis. *Science.* 2000; 289:920-930.
6. Buskirk AR, Green R. Ribosome pausing, arrest and rescue in bacteria and eukaryotes. *Philosophical Transactions of the Royal Society B-Biological Sciences.* 2017;372.
7. Schuller AP, Green R. Roadblocks and resolutions in eukaryotic translation. *Nature Reviews Molecular Cell Biology.* 2018;19:526-541.
8. Youngman EA, McDonald ME, Green R. Peptide Release on the Ribosome: Mechanism and Implications for Translational Control. *Annual Review of Microbiology.* 2008;62:353-373.
9. Zaher HS, Green R. Fidelity at the Molecular Level: Lessons from Protein Synthesis. *Cell.* 2009;136:746-762.
10. Pavlov MY, Liljas A, Ehrenberg M. A recent intermezzo at the Ribosome Club. *Philosophical Transactions of the Royal Society B-Biological Sciences.* 2017;372.
11. Raine A, Lovmar M, Wikberg J, Ehrenberg M. Trigger factor binding to ribosomes with nascent peptide chains of varying lengths and sequences. *Journal of Biological Chemistry.* 2006;281:28033-28038.
12. Ullers RS, Houben EN, Raine A, ten Hagen-Jongman CM, Ehrenberg M, Brunner J, et al. Interplay of signal recognition particle and trigger factor at L23 near the nascent chain exit site on the Escherichia coli ribosome. *Journal of Cell Biology.* 2003;161:679-684.
13. Chaney JL, Clark PL. Roles for Synonymous Codon Usage in Protein Biogenesis. *Annual Review of Biophysics.* 2015;44:143-166.
14. Liu Y, Yang Q, Zhao FZ. Synonymous but Not Silent: The Codon Usage Code for Gene Expression and Protein Folding. *Annual Review of Biochemistry.* 2021;90:375-401.
15. Nissley DA, O'Brien EP. Timing Is Everything: Unifying Codon Translation Rates and Nascent Proteome Behavior. *Journal of the American Chemical Society.* 2014;136:17892-17898.
16. Czech A, Fedyunin I, Zhang G, Ignatova Z. Silent mutations in sight: co-variations in tRNA abundance as a key to unravel consequences of silent mutations. *Molecular Biosystems.* 2010;6:1767-1772.
17. Rauscher R, Ignatova Z. Timing during translation matters: synonymous mutations in human pathologies influence protein folding and function. *Biochemical Society Transactions.* 2018;46:937-944.

Citation: Guzman-Luna V, Cavagnero S. Mg²⁺ ions mediate the interaction of intrinsically disordered nascent chains with the ribosome: implications for protein folding and aggregation in the early stages of protein life. *Neurosci Chron.* 2022;3(1):3-5.

18. Zhang G, Ignatova Z. Folding at the birth of the nascent chain: coordinating translation with co-translational folding. *Current Opinion in Structural Biology.* 2011;21:25-31.
19. Evans MS, Sander IM, Clark PL. Cotranslational Folding Promotes beta-Helix Formation and Avoids Aggregation In Vivo. *Journal of Molecular Biology.* 2008;383:683-692.
20. Addabbo RM, Dalphin MD, Mecha MF, Liu Y, Staikos A, Guzman-Luna V, et al. Complementary Role of Co- and Post-Translational Events in De Novo Protein Biogenesis. *Journal of Physical Chemistry B.* 2020;124:6488-6507.
21. Cruz-Vera LR, Rajagopal S, Squires C, Yanofsky C. Features of ribosome-peptidyl-tRNA interactions essential for tryptophan induction of trn operon expression. *Molecular Cell.* 2005;19:333-343.
22. Rodnina MV, Wintermeyer W, Green R. Ribosomes : structure, function, and dynamics. New York: SpringerWien; 2011.
23. Seidelt B, Innis CA, Wilson DN, Gartmann M, Armache JP, Villa E, et al. Structural insight into nascent polypeptide chain-mediated translational stalling. *Science.* 2009;326:1412-1415.
24. Bhushan S, Hoffmann T, Seidelt B, Frauenfeld J, Mielke T, Berninghausen O, et al. SecM-stalled ribosomes adopt an altered geometry at the peptidyl transferase center. *PLoS Biology.* 2011;9:e1000581.
25. Orfanoudaki G, Economou A. Proteome-wide subcellular topologies of E. coli polypeptides database (STEPdb). *Mol Cell Proteomics.* 2014;13:3674-3687.
26. Knight AM, Culviner PH, Kurt-Yilmaz N, Zou T, Ozkan SB, Cavagnero S. Electrostatic Effect of the Ribosomal Surface on Nascent Polypeptide Dynamics. *ACS Chemical Biology.* 2013;8:1195-1204.
27. BurrIDGE C, Waudby CA, Wlodarski T, Cassaignau AME, Cabrera LD, Christodoulou J. Nascent chain dynamics and ribosome interactions within folded ribosome-nascent chain complexes observed by NMR spectroscopy. *Chemical Science.* 2021;12:13120-13126.
28. Cassaignau AME, Wlodarski T, Chan SHS, Woodburn LF, Bukvin IV, Streit JO, et al. Interactions between nascent proteins and the ribosome surface inhibit co-translational folding. *Nature Chemistry.* 2021;13:1214.
29. Deckert A, Waudby CA, Wlodarski T, Wentink AS, Wang XL, Kirkpatrick JP, et al. Structural characterization of the interaction of alpha-synuclein nascent chains with the ribosomal surface and trigger factor. *Proceedings of the National Academy of Sciences of the United States of America.* 2016;113:5012-5017.
30. Deckert A, Cassaignau AME, Wang XL, Wlodarski T, Chan SHS, Waudby CA, et al. Common sequence motifs of nascent chains engage the ribosome surface and trigger factor. *Proceedings of the National Academy of Sciences of the United States of America.* 2021;118.
31. Guzman-Luna V, Fuchs AM, Allen AJ, Staikos A, Cavagnero S. An intrinsically disordered nascent protein interacts with specific regions of the ribosomal surface near the exit tunnel. *Communications Biology.* 2021;4.
32. Kolev SK, Petkov PS, Rangelov MA, Trifonov DV, Milenov TI, Vayssilov GN. Interaction of Na(+), K(+), Mg(2+) and Ca(2+) counter cations with RNA. *Metallomics.* 2018;10:659-678.
33. Nierhaus KH. Mg²⁺, K⁺, and the Ribosome. *Journal of Bacteriology.* 2014;196:3817-3819.
34. Southall NT, Dill KA, Haymet ADJ. A view of the hydrophobic effect. *Journal of Physical Chemistry B.* 2002;106:521-533.
35. Record MT, Courtenay ES, Cayley DS, Guttman HJ. Responses of E-coli to osmotic stress: Large changes in amounts of cytoplasmic solutes and water. *Trends in Biochemical Sciences.* 1998;23:143-148.
36. Record MT, Courtenay ES, Cayley S, Guttman HJ. Biophysical compensation mechanisms buffering E-coli protein-nucleic acid interactions against changing environments. *Trends in Biochemical Sciences.* 1998;23:190-194.
37. Wendel BM, Pi HL, Kruger L, Herzberg C, Stulke J, Helmann JD. A Central Role for Magnesium Homeostasis during Adaptation to Osmotic Stress. *Mbio.* 2022;13.
38. Yoshimune K, Yoshimura T, Esaki N. Hsc62, a new DnaK homologue of Escherichia coli. *Biochemical and Biophysical Research Communications.* 1998;250:115-118.
39. Sriram M, Osipiuk J, Freeman BC, Morimoto RI, Joachimiak A. Human Hsp70 molecular chaperone binds two calcium ions within the ATPase domain. *Structure.* 1997;5:403-414.
40. Yeom J, Groisman EA. Low Cytoplasmic Magnesium Increases the Specificity of the Lon and ClpAP Proteases. *Journal of Bacteriology.* 2021;203.
41. Iotti S, Malucelli E. In vivo assessment of Mg(2+) in human brain and skeletal muscle by (31)P-MRS. *Magnesium Research.* 2008;21:157-162.
42. Sun Y, Cruz AM, Hadley KC, Galant NJ, Law R, Vernon RM, et al. Physiologically Important Electrolytes as Regulators of TDP-43 Aggregation and Droplet-Phase Behavior. *Biochemistry.* 2019;58:590-607.
43. Darling AL, Uversky VN. Intrinsic Disorder in Proteins with Pathogenic Repeat Expansions. *Molecules.* 2017;22.
44. Uversky VN. Intrinsic disorder in proteins associated with neurodegenerative diseases. *Frontiers in Bioscience-Landmark.* 2009;14:5188-5238.
45. Duc KD, Batra SS, Bhattacharya N, Cate JHD, Song YS. Differences in the path to exit the ribosome across the three domains of life. *Nucleic Acids Research.* 2019;47:4198-4210.
46. de Jesus V, Qureshi NS, Warhaut S, Bains JK, Dietz MS, Heilemann M, et al. Switching at the ribosome: riboswitches need rProteins as modulators to regulate translation. *Nature Communications.* 2021;12.
47. Singh R, Rao CM. Chaperone-like activity and surface hydrophobicity of 70S ribosome. *FEBS Letters.* 2002;527:234-238.
48. Deuerling E, Gamberdinger M, Kreft SG. Chaperone Interactions at the Ribosome. *Cold Spring Harbor Perspectives in Biology.* 2019;11.